

FcCH₂P(CH₂OH)₂: a new, reactive yet air-stable ferrocene-derived phosphine [Fc = (η-C₅H₅)FeC₅H₄]

Nicholas J. Goodwin, William Henderson* and J. Kwabena Sarfo

Department of Chemistry, University of Waikato, Private Bag 3105, Hamilton, New Zealand

Reaction of FcCH₂NMe₃⁺I⁻ with an excess of P(CH₂OH)₃ gives the air-stable ferrocenyolphosphine FcCH₂P(CH₂OH)₂, characterised by an X-ray structure determination; reactivity towards reagents such as MeI, acrylonitrile and amines allows the synthesis of a range of new ferrocene-phosphine derivatives.

The chemistry of phosphine ligands containing the ferrocenyl group has attracted much interest over a period of many years, and there is an extensive literature on the topic.¹ Such phosphines are of interest for their coordination chemistry, including their applications to asymmetric catalysis.² In contrast to the large number of ferrocene-derived phosphines, and their metal complexes, which have the phosphorus directly bonded to a cyclopentadienyl ring, ligands where there is a carbon spacer between these two functionalities are somewhat rarer,³ though chiral versions are attracting interest for their catalytic activity.⁴ We are interested in developing the chemistry of hydroxymethylphosphines which also contain the ferrocenyl group. Hydroxymethylphosphines contain chemically reactive PCH₂OH groups,⁵ allowing further derivatisation of the phosphine ligand. Examples include the synthesis [from P(CH₂OH)₃] of ligands such as P(CH₂CH₂CN)₃⁶ and the water-soluble phosphotriazaadamantane, PTA **1**,⁷ both of which have attracted recent attention. In addition, hydroxymethylphosphines themselves have been the subjects of a number of recent studies, for both their water-solubility and reactivity.⁸ Here we report the first example of a ferrocene-derived hydroxymethylphosphine ligand, together with some derivatives.

Reaction of the readily available ferrocenylmethyl ammonium salt FcCH₂NMe₃⁺I⁻⁹ with an excess of P(CH₂OH)₃[†] in refluxing methanol followed by treatment with an excess of triethylamine, and extraction with diethyl ether gave orange crystals of FcCH₂P(CH₂OH)₂ **2**, as shown in Scheme 1. Recrystallisation from dichloromethane–methanol–light petroleum (bp 40–60 °C) gave the product in 66% overall yield. The compound is air-stable, both in the solid state and in solution; other hydroxymethylphosphines are typically air-sensitive to some extent.

The formulation of **2** was confirmed by elemental analysis, NMR spectroscopy and electrospray mass spectrometry,[‡] together with an X-ray structure determination.§ The molecular structure is shown in Fig. 1, together with selected bond lengths and angles. The structure determination of **2** is the first of a free

hydroxymethylphosphine. The molecule forms a hydrogen-bonded array, propagated along the 2₁ screw axis, parallel to the *a*-axis of the unit cell; the resulting ten-membered rings are formed by hydrogen bonding between three molecules of **2**.

Treatment of **2** with hydrogen peroxide or elemental sulfur afforded the phosphine oxide (δ_P 45.8, CDCl₃) and sulfide (δ_P 47.6, CDCl₃) derivatives FcCH₂P(X)(CH₂OH)₂ (X = O, S) respectively. Representative metal complexes of **2**, *cis*-[PtCl₂(**2**)₂] and [AuCl(**2**)] were readily prepared by reaction of a dichloromethane solution of **2** with [PtCl₂(cod)] (cod = cycloocta-1,5-diene) and [AuCl(CNBU[†])] respectively. The ³¹P NMR spectrum of the platinum complex showed the expected single resonance, with ¹⁹⁵Pt satellites [*J*(PtP) 3427 Hz].

The unique reactivity of hydroxymethylphosphines facilitates the ready synthesis of other functionalised phosphines derived from **2**. Thus, reaction of **2** with an excess of MeI yields the phosphonium salt FcCH₂P(Me)(CH₂OH)₂⁺I⁻ **3**, characterised by an X-ray structure determination and electrospray mass spectrometry (*m/z* 307, *M*⁺). Subsequent treatment of **3** with an excess of triethylamine generates the mono(hydroxymethyl)phosphine FcCH₂P(Me)CH₂OH (δ_P -33.1, CDCl₃). Reaction of **2** with an excess of acrylonitrile affords FcCH₂P(CH₂CH₂CN)₂ (δ_P -22.1, CDCl₃), while reaction with an excess of diethylamine proceeds rapidly *via* a Mannich-type condensation reaction,⁵ giving FcCH₂P(CH₂NEt₂)₂, characterised as the phosphine oxide (δ_P 45.0, CDCl₃). This reactivity towards amine groups can be exploited to immobilise the phosphine **2** on amine-containing supports, such as 3-aminopropylsilica. The phosphine **2** thus represents a convenient new reagent for the incorporation of ferrocene groups into amine-containing materials.





Fig. 1 Molecular structure of FcCH₂P(CH₂OH)₂ **2**, with atom numbering scheme. Selected bond lengths (Å) and angles (°) C(1)–C(11) 1.489(6), P–C(1) 1.858(4), P–C(2) 1.862(4), P–C(3) 1.851(4), C(2)–O(2) 1.429(5), C(3)–O(3) 1.423(5), Fe–C 2.034(4)–2.047(5), average 2.040(4), C(11)–C(1)–P 114.6(3), C(1)–P–C(2) 95.0(2), C(1)–P–C(3) 101.3(2), C(2)–P–C(3) 99.5(2), P–C(2)–O(2) 111.9(3), P–C(3)–O(3) 116.2(3).



We thank the University of Waikato and the New Zealand Lottery Grants Board for financial support, and Professor Ward Robinson (University of Canterbury, New Zealand) for acquisition of the X-ray data sets. N. J. G. thanks the William Georgetti Trust for a scholarship, and Professor B. K. Nicholson for assistance with the crystallography.

Footnotes

† $\text{P}(\text{CH}_2\text{OH})_3$ was prepared by the reaction of methanolic $\text{P}(\text{CH}_2\text{OH})_4\text{Cl}$ (obtained as Retardol C from Albright & Wilson Ltd., UK, as an 80% m/m aqueous solution) with 1 mol equiv. of KOH under a nitrogen atmosphere.

‡ Anal. Found: C, 52.9; H, 6.0. $\text{C}_{13}\text{H}_{17}\text{FeO}_2\text{P}$ requires C, 53.5; H, 5.9%. Mp 104–106 °C. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3), δ –19.3. Electrospray mass spectrometric analysis with added AgNO_3 ¹⁰ at a cone voltage of 60 V showed the following ions (m/z): $[2\text{M} + \text{Ag}]^+$ (690.8), $[\text{M} + \text{Ag}]^+$ (398.9), $[\text{M}]^+$ (292.1) and $[\text{FcCH}_2]^+$ (199.0). The observation of the ferrocenium ion may be as a result of oxidation by silver ions, or alternatively by electrochemical oxidation in the electrospray capillary.¹¹

§ Accurate cell parameters and intensity data were recorded on a Nicolet R3 diffractometer, using a crystal of dimensions $0.70 \times 0.45 \times 0.25$ mm, at –115 °C. Crystal data: $M_r = 292.10$; orthorhombic, space group $P2_12_12_1$, $a = 6.299(2)$, $b = 11.561(2)$, $c = 17.519(2)$ Å, $U = 1275.8(4)$ Å³, $D_c = 1.521$ g cm^{–3} ($Z = 4$), $F(000) = 608$, $\mu(\text{Mo-K}\alpha) = 1.29$ mm^{–1}, $\lambda = 0.71073$ Å. A total of 1421 reflections were collected using ω -scans in the range $4 < 2\theta < 50^\circ$. Of these 1401 were unique ($R_{\text{merge}} = 0.0140$ after the data were corrected for absorption based on a series of psi-scans). The structure was solved by direct methods and developed routinely. Full-matrix least-squares refinement was based on F^2 , with all non-hydrogen atoms anisotropic and with ring and CH_2 hydrogen atoms included in calculated positions with isotropic temperature factors 1.2 times that of the U_{iso} of the atom to which they are bonded. The hydroxy hydrogen atoms were located in a penultimate electron density map and were refined with fixed isotropic temperature factors. The refinement converged at $R_1 = 0.0297$ [for 1242 reflections with $I > 2\sigma(I)$] and $wR_2 = 0.0667$ (all data) [$w = [\sigma^2(F_o^2) + (0.0593P)^2]^{-1}$ where $P = (F_o^2 + 2F_c^2)/3$]; final GOF = 0.957. The final difference map showed no peak $> +0.385$ e Å^{–3} and no hole < -0.294 e Å^{–3}. A Flack x parameter of 0.04(3) showed the refinement had been carried out with the correct polarity. Programs used were SHELXS-86¹² and SHELXL-93.¹³ Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre (CCDC). See Information for Authors, Issue No. 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 182/101.

References

- 1 For selected recent references, see: K.-S. Gan and T. S. A. Hor, *Ferrocenes*, ed. A. Togni and T. Hayashi, VCH, Weinheim, 1995, p. 1; I. R. Butler, M. Kalaji, L. Nehrlich, M. Hursthouse, A. I. Karaulov and K. M. A. Malik, *J. Chem. Soc., Chem. Commun.*, 1995, 459.
- 2 A. Mernyi, C. Kratky, W. Weissensteiner and M. Widhalm, *J. Organomet. Chem.*, 1996, **508**, 209; J. M. Brown, *Chem. Soc. Rev.*, 1993, 25; O. Reiser, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 547; A. Yamazaki, T. Morimoto and K. Achiwa, *Tetrahedron: Asymmetry*, 1993, **4**, 2287; M. Sawamura and Y. Ito, *Chem. Rev.*, 1992, **92**, 857.
- 3 See for example: S. Hoppe, H. Weichmann, K. Jurkschat, C. Schneider-Koglin and M. Dräger, *J. Organomet. Chem.*, 1995, **505**, 63; Y. Yamamoto, T. Tanase, I. Mori and Y. Nakamura, *J. Chem. Soc., Dalton Trans.*, 1994, 3191.
- 4 Recent references: N. C. Zanetti, F. Spindler, J. Spencer, A. Togni and G. Rihs, *Organometallics*, 1996, **15**, 860; P. Barbaro and A. Togni, *Organometallics*, 1995, **14**, 3570; H. C. L. Abbenhuis, U. Burckhardt, V. Gramlich, C. Köllner, P. S. Pregosin, R. Salzmann and A. Togni, *Organometallics*, 1995, **14**, 759; M. Sawamura, H. Hamashima, M. Sugawara, R. Kuwano and Y. Ito, *Organometallics*, 1995, **14**, 4549; M. Sawamura, R. Kuwano and Y. Ito, *J. Am. Chem. Soc.*, 1995, **117**, 9602.
- 5 A. W. Frank, D. J. Daigle and S. L. Vail, *Textile Res. J.*, 1982, 738.
- 6 C. W. Liu, H. Pan, J. P. Fackler, Jr., G. Wu, R. E. Wasylischen and M. Shang, *J. Chem. Soc., Dalton Trans.*, 1995, 3691; M. N. I. Khan, C. King, J. P. Fackler, Jr. and R. E. Wimpenny, *Inorg. Chem.*, 1993, **32**, 2502.
- 7 J. M. Forward, Z. Assefa, R. J. Staples and J. P. Fackler, Jr., *Inorg. Chem.*, 1996, **35**, 16; J. M. Forward, D. Bohmann, J. P. Fackler, Jr. and T. J. Staples, *Inorg. Chem.*, 1995, **34**, 6330; B. Assmann, K. Angermaier, M. Paul, J. Riede and H. Schmidbaur, *Chem. Ber.*, 1995, **128**, 891; D. J. Darensbourg, N. W. Stafford, F. Joó and J. H. Reibenspies, *J. Organomet. Chem.*, 1995, **488**, 99; Z. Assefa, B. G. McBurnett, R. J. Staples, J. P. Fackler, Jr., B. Assmann, K. Angermaier and H. Schmidbaur, *Inorg. Chem.*, 1995, **34**, 75; J. M. Forward, J. P. Fackler, Jr. and R. J. Staples, *Organometallics*, 1995, **14**, 4191; Z. Assefa, B. G. McBurnett, R. J. Staples and J. P. Fackler, Jr., *Inorg. Chem.*, 1995, **34**, 4965; B. Assmann, K. Angermaier and H. Schmidbaur, *J. Chem. Soc., Chem. Commun.*, 1994, 941.
- 8 M. Slany, M. Bardajf, M.-J. Casanove, A.-M. Caminade, J.-P. Majoral and B. Chaudret, *J. Am. Chem. Soc.*, 1995, **117**, 9764; F. Bitterer, F. Kucken and O. Stelzer, *Chem. Ber.*, 1995, **128**, 275; H. H. Petach, W. Henderson and G. M. Olsen, *J. Chem. Soc., Chem. Commun.*, 1994, 2181; J. Fawcett, P. A. T. Hoye, R. D. W. Kemmitt, D. J. Law and D. R. Russell, *J. Chem. Soc., Dalton Trans.*, 1993, 2563; J. W. Ellis, K. N. Harrison, P. A. T. Hoye, A. G. Orpen, P. G. Pringle and M. B. Smith, *Inorg. Chem.*, 1992, **31**, 3026; *Platinum Met. Rev.*, 1990, **34**, 74.
- 9 D. Lednicer and C. R. Hauser, *Org. Synth. Coll.*, 1973, **5**, 434.
- 10 W. Henderson and G. M. Olsen, *Polyhedron*, 1996, **15**, 2105.
- 11 W. Henderson, A. G. Oliver and A. L. Downard, *Polyhedron*, 1996, **15**, 1165; X. Xu, S. P. Nolan and R. B. Cole, *Anal. Chem.*, 1994, **66**, 119.
- 12 G. M. Sheldrick, SHELXS-86, Program for solving crystal structures, University of Göttingen, 1986.
- 13 G. M. Sheldrick, Program for refining crystal structures, University of Göttingen, 1993.

Received, 9th April 1996; Com. 6/02420C