

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331666339>

Comparative Analysis of Haematological Parameters in Hookworm and Plasmodium falciparum Co-Infected Individuals in Kintampo North Municipality, Ghana

Article in *International Journal of TROPICAL DISEASE & Health* · March 2019

DOI: 10.9734/ijtdh/2018/v3i4i430101

CITATION

1

READS

166

7 authors, including:



Benjamin Amoani

University of Cape Coast

16 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)



Ben Gyan

Noguchi Memorial Institute for Medical Research

130 PUBLICATIONS 1,193 CITATIONS

[SEE PROFILE](#)



Francis Armah

University of Cape Coast

23 PUBLICATIONS 40 CITATIONS

[SEE PROFILE](#)



Samuel Kofi Tchum

Kintampo Health Research Centre

30 PUBLICATIONS 268 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Analytical method development and validation for research [View project](#)



Mentorship [View project](#)



Comparative Analysis of Haematological Parameters in Hookworm and *Plasmodium falciparum* Co-Infected Individuals in Kintampo North Municipality, Ghana

**Benjamin Amoani^{1,2,3*}, Ben Gyan^{2*}, Francis A. Armah¹, Cecil Otabil¹,
Kofi Tchum⁴, Margaret T. Frempong³ and Michael D. Wilson⁵**

¹Department of Biomedical Science, School of Allied Health Sciences, University of Cape Coast, Cape Coast, Ghana.

²Department of Immunology, Noguchi Memorial Institute for Medical Research, College of Health Sciences, University of Ghana, Legon, Ghana.

³Department of Molecular Medicine, School of Medical Sciences, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

⁴Kintampo Health Research Centre, Ghana Health Service, Ministry of Health, Kintampo, Ghana.

⁵Department of Parasitology, Noguchi Memorial Institute for Medical Research, College of Health Sciences, University of Ghana, Legon, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2018/v34i430101

Editor(s):

(1) Dr. Giuseppe Murdaca, Clinical Immunology Unit, Department of Internal Medicine, University of Genoa, Italy.

Reviewers:

(1) Loick Pradel Kojom, The University of Douala, Cameroon.

(2) Franco Cervellati, University of Ferrara, Italy.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47535>

Received 11 November 2018

Accepted 22 February 2019

Published 11 March 2019

Original Research Article

ABSTRACT

Background: Hookworm and *Plasmodium falciparum* are endemic in Ghana, especially in the north-east and middle belt areas. Haematological parameters have been shown to predict the presence of these infections in patients. This study investigated the effect of hookworm-*P. falciparum* co-infection on haematological parameters among the patients.

Methodology: Stool and blood samples from 984 participants in a cross-sectional study conducted

*Corresponding author: Email: bamoani@ucc.edu.gh, bgyan@noguchi.ug.edu.gh;

in the Kintampo North Municipality of Ghana were examined for hookworm and *Plasmodium falciparum* parasites. Malaria parasitaemia was estimated by microscopy and *Plasmodium falciparum*-specific 18S rRNA gene by polymerase chain reaction. Hookworm eggs in faecal samples were analyzed using Kato-Katz and formol-ether concentration methods. Hookworm speciation was done by PCR. Estimation of haematological parameters was done by automated haematology analyzer. Tukey multiple comparisons test was used to compare continuous variables among the infected groups and Spearman's rank correlation test determined the relationships between variables.

Results: The overall mean (SD) age of the study participants was 22.8 (17.4). Mean lymphocytes and eosinophils counts were higher while mean neutrophil and monocytes counts were lower in the co-infected individuals compared to the single infections. The mean red blood cell (RBC) count and haemoglobin (Hb) levels were higher in the co-infected individuals than in those with malaria only infection. However, white blood cell count and platelet counts were not statistically significant among the groups. There was a significant negative correlation between malaria parasite density and Hb level ($r = -0.239$, $p = 0.001$)

Conclusion: Hookworm and *Plasmodium falciparum* co-infections showed an increased in lymphocyte, eosinophil count and Haemoglobin levels, but a decrease in neutrophil and monocyte levels compared to malaria only infection. The possible mechanisms accounting for the variations in haematological levels remain to be elucidated.

Keywords: Hookworm; *plasmodium falciparum*; parasitaemia; infection intensity; haematological parameter.

1. INTRODUCTION

Hookworm is estimated to affect about 740 million people in the world [1], including 156 million children and most of these individuals are found in tropical regions of the world where such infections are linked to poverty [2]. In sub-Saharan Africa, hookworm prevalence is approximately 30 % [3]. However, in the north-eastern Ghana and the middle belt of Ghana (Kintampo North Municipality), the prevalence has been reported to be as high as 50 % [4] and 45 % [5]. The resultant effects of hookworm infection include growth delay, malnutrition, poor appetite and anemia, which, in pregnancy may result to poor birth outcomes [6]. Hookworm infection may also cause retardation in both physical and cognitive development in young children [7,8].

Malaria is one of the leading causes of morbidity and mortality in the developing world, especially sub-Saharan Africa. In endemic areas, about 60–70 % of the cases are attributable to *Plasmodium falciparum* infection while 30–40 % are attributable to other malaria parasite infections [9,10]. *P. falciparum* is responsible for 13–28 % of deaths in children under 5 years of age [11]. The high prevalence of both malaria and hookworm infections among individuals living in Africa means that a co-infection will be common [12]. However, little is known about the interaction between these widely distributed parasites. Hookworms cause chronic intestinal

blood loss while acute haemolysis and depletion of haemoglobin are associated with *Plasmodium* infections [13-14]. Therefore, there is the need to investigate the haematological profiling in hookworm and malaria co-infection. Unfortunately, not enough studies have been done to investigate the pathological effects of *N. americanus* and *P. falciparum* infections to determine how their co-occurrence, as well as individual occurrences, may affect an individual's general blood cell levels. Results from studies addressing this effect would help to possibly predict the type as well as the level of infection, with respect to hookworm and malaria, which would go a long way to have profound implications for both malaria and hookworm control programmes in Ghana.

2. METHODS

2.1 Study Site, Design and Recruitment of Participants

The study was approved by the Institutional Review Board of Noguchi Memorial Institute for Medical Research (FWA#: 00001824). All study participants provided written informed consent prior to their recruitment. This study was conducted in Kintampo North Municipality located within the forest-savannah transitional ecological zone in the middle belt of Ghana. The ages of the study participants ranges from 4yrs to 80 yrs.

2.2 Sample Collection and Processing

Trained field workers administered demographic and health questionnaires, and distributed labeled stool-collection containers to the participants. Stool samples were collected the following day and finger pricks were made to test for malaria using Rapid Diagnostic Test (RDT) kits (CareStart™ Malaria PfHRP2/pLDH Ag RDT, Access Bio, Inc, USA) and to prepare thin and thick blood film on the same slide. About 5 mL of blood was drawn at the same time into EDTA vacutainers tubes for haematological analysis. Separate samples of blood were spotted on Whatman FTA Blood Stain Cards for storage until use in species identification using PCR. Prepared slides were stained with Giemsa and examined under the light microscope. Malaria parasitaemia was estimated by microscopy according to WHO protocols [15] and *P. falciparum*-specific 18S rRNA gene was detected in blood by PCR. Faecal samples were analyzed for the presence of helminth eggs on the day of collection using the Kato-Katz and formol-ether concentration methods. Hookworm speciation was carried out for hookworm positive cases by PCR using specific primers.

2.3 Hookworm Speciation by PCR

Hookworm species identification was determined using genomic DNA extracted from purified hookworm eggs samples of infected individuals using QIAamp DNA stool kit (QIAGEN, Hilden, Germany). Five microliters of purified gDNA (20-40 ng) was amplified in 1.25 mM each of deoxynucleotide triphosphate (dNTP), 1U of the *Taq* DNA polymerase enzyme (Sigma, Cat. #. D1806-250UN) and 0.3µL of each primer. The primers used were forward primer (NC2; 5'-TTA GTT TCT TTT CCT CCG CT-3'), with species specific reverse primers for *A. duodenale* (jmAD; 5'-TGC GAA GTT CGC GTT CGC TGA GC-3') or *N. americanus* (jmNA; 5'-CGT TAA CAT TGT ATA CCT GTA CAT AC-3') in separate reactions as described elsewhere [16]. The amplification conditions were initial heating at 94°C for 5 minutes, followed by 40 cycles of denaturation at 94°C for 1 minute, annealing at 55°C for 1 minute and extension at 72°C for 1 minute, with a final elongation step at 72°C for 5 minutes.

2.4 PCR Identification of *P. falciparum*

Total DNA was extracted from FTA cards using the Chelex method [17]. A 276 bp fragment of *P. falciparum* 18S rRNA gene sequence was amplified using the specific forward 5'-AAC AGA

CGG GTA GTC ATG ATT GAG-3' and reverse 5'-GTA TCT GAT CGT CTT CAC TCCC-3' primers as used elsewhere [18]. The 20 µl reaction contained 20 – 40 ng total DNA, 0.25 mM of each primer, 1.25 mM of each dNTP, 1U of HotStar *Taq*® DNA polymerase (Biomol GmbH, Hamburg, Germany) and 1X reaction buffer. The PCR conditions were 34 cycles of denaturation at 94 °C for 30 seconds, annealing at 54°C for 30 seconds and extension at 72 °C for 1 minute with a final elongation step at 72 °C for 5 minutes. DNA of the NF54 strain of *P. falciparum* extracted from culture was included on each PCR plate as positive control.

2.5 Visualization of Amplicons

The amplified products were visualized and the sizes determined by UV visualization after electrophoresis in a 2 % ethidium bromide stained-agarose gel. Products of the appropriate size (690 bp for *A. duodenale*, 870 bp for *N. americanus* and 276 bp for *P. falciparum*) were considered positive compared to standard controls.

2.6 Haematological Profiling

The haematological levels were determined using haematology analyser (ABX Pentra 60C+, HORIBA Medical, Rue du Caduce', France) by following the manufacturer's instructions.

2.7 Statistical Analysis

Statistical analysis was done by SPSS Version 24 (Chicago, IL, USA). Proportions such as prevalence were compared between groups (Pearson χ^2 test). Normalized variables were compared between groups using either the Welch two sample t test or One-Way analysis of variance (ANOVA) where appropriate. Pairwise differences between groups were compared using Post Hoc Test (Turkey's HSD). Spearman's rank correlation test determined the relationships between variables. P-value of <0.05 was considered statistically significant.

3. RESULTS

3.1 Demographic and Parasitological Characteristics of the Study Population

Table 1 shows the demographic characteristics of the study population. The study population consisted of 48.9 % males and 51.1 % females. The overall mean (SD) age of the study participants was 22.8 (17.4). There was no

significant difference in the geometric mean of egg per gram (EPG) of stool for individuals who were infected with only hookworm and those with *P.falciparum* and hookworm co-infection, though the co-infected group showed a marginal increase in intensity of hookworm infection. The mean *P. falciparum* density of individuals with malaria infection mono-infection was also significantly higher than those individuals co-infected with hookworm ($p=0.001$). All the PCR speciation for hookworm showed positive for *Necator americanus*.

3.2 Relationship between Haematological Parameters and Infection Status

Significant changes were found in the haematological parameters of the control, healthy subjects and the hookworm and malaria infected individuals. Mean white blood cell (WBC) counts did not generally vary significantly among the various infection statuses ($p = 0.056$) even though, within the various infected groups, the mean WBC level was significantly higher in the malaria-mono infected group than in the hookworm-mono infected group ($p = 0.0348$) which had the least mean WBC count (Table 2). Mean neutrophil, monocyte, eosinophil, lymphocyte, red blood cell (RBC) and haemoglobin levels all exhibited a significant variations among the various groups generally. It was also observed that co-infection caused an increase in lymphocyte, eosinophil, red blood cell, and haemoglobin levels compared to the malaria only. However, the co-infection led to significant reduction in neutrophil and monocyte level compared to malaria-mono infected group. Basophil levels were found to be increased in infected individuals with the highest level found to be associated with the hookworm-only infected individuals. Platelet counts, showed no significant variations among the groups.

3.3 Association of Intensity of Infection with Haematological Parameter

Table 3 shows the relationship between laboratory parameters and intensity of infection in individuals with hookworm and malaria infections. Lymphocytes, neutrophil and basophil levels correlated negatively, and weak ($r = -0.020$; $r = -0.103$; $r = -0.017$) with *P. falciparum* intensity and statistically not significant. But, correlation between *P. falciparum* intensity and haemoglobin (Hb) levels was negative, medium and statistically significant ($r = -0.237$, $p=0.001$). The relationship of *P. falciparum* intensity with relative eosinophil count and monocytes showed

medium, positive ($r = 0.281$, $p=0.036$; $r = 0.154$, $p<0.001$) and a statistically significant correlation. The relationship of *N. americanus* intensity with WBC, neutrophils and monocytes showed a medium and strong, negative respectively ($r = -0.235$; $r = -0.437$; -0.562) and a statistically highly significant correlation. Though there was also a negative correlation between *N. americanus* intensity and Hb, it was not significant.

4. DISCUSSION

Complete blood count is a routine haematological test frequently used to help diagnose a number of diseases, such as anaemia, various acute infections, immune disorders, cancers and in health screening [19]. Hookworm infection and malaria are both known to cause anemia [20, 21]. Both infections will characteristically induce immune responses in the body like all other infections. However, due to differences in the anatomical position of the parasites and the mechanism of feeding or infecting of RBCs within the host. Hookworm harbours in the small intestine to obtain its food, thus lives outside the body cells, whilst *P. falciparum* infects the hepatic and the RBCs, thus lives within the cells. Therefore, different specific immune responses mediated by specific immune cells are expected to be elicited towards these infections.

In this study, significant changes in RBC counts and haemoglobin levels were found with malaria-mono infected subjects having the least RBC and haemoglobin levels compared to the co-infected group. We observed that increased in parasite density led to a significant reduction in Hb level. The destruction of RBCs by malaria parasite rapid proliferation and clearance of malaria-infected RBCs by the immune system are contributory factors to the severity of malarial anaemia [22,23]. Hookworm infection on the other hand, results in only intestinal blood loss, where the parasite resides and feeds on blood, as the cause of anemia [24]. Therefore, we expected co-infections to experience reduced Hb and RBCs level [25]. However, our result showed that the levels of RBCs and Hb were higher in the co-infected individual than the malaria-mono infected individuals, thus, suggesting some protective effect in the presence of co-infection [4,26,27]. The mechanism by which hookworm apparently protect against a decrease in haemoglobin in *P. falciparum* malaria is unknown. The high levels of the Th2 cytokines (IL-10) produce during helminth infection may

counteract the Th1 cytokines (TNF-alpha) induced by malaria to prevent the development of severe anemia [28]. The overall Th2/Th1 balance, the homeostatic role of interleukin 10 and TGF-β as modulators of the immune response [29], and the role of the CD23/NO pathway in reducing sequestration [30] are additional possible mechanisms of protection against severe malaria [27].

It was also found that lymphocyte numbers were significantly increased in the co-infected individuals compared to malaria only infected individual, however, similar to those with hookworm mono-infection. The reason for this may be due to the complexity of the hookworm life-cycle which offers numerous opportunities for parasite-host interaction at the molecular level. Additionally, natural attrition of larvae at critical barriers, such as during skin invasion, and transit through lung tissues, as well as arrival in the gut and penetration of its mucosa, presents the host with an extensive diversity of antigenic challenge, immune stimulation and modulation [21]. This is opposed to the relatively simple life-cycle of the malaria parasites which only involves the hepatic and erythrocytic stages in the host.

Neutropenia was observed in individuals with co-infection compared to malaria only infected individuals. This possibly may be due to a decrease in production, increased destruction or an accelerated usage of neutrophils which usually occurs during concurrent infections. Secreted proteins by the infective larvae of *N. americanus* (*Na*-ASP-2) has been found to induce significant leukocyte (mostly comprised of 60% neutrophils and 30% monocytes) influx to the skin [31]. Furthermore, Neutrophil Inhibitory Factor (NIF), a glycoprotein secreted by the adult *N. americanus* may also be a possible contributor to the low neutrophil numbers due to its ability to potently inhibit CD11b/CD18-dependent neutrophil function and recruiting at worm attachment sites [32, 33]. Also, other factors such as vitamin B12 deficiency or unmeasured infections may have accounted for

this low neutrophil count as these factors are known to negatively impact neutrophil levels [34]. Monocyte counts was significantly found to be higher in malaria only infected individuals compared to those with co-infection or single infected with hookworm, which corroborate with other studies [35,36]. Mononuclear cells, which are activated by *Plasmodium* during malarial infection, produce inflammatory cytokines such as tumour necrosis factor alpha (TNF-α), interleukin-1 (IL-1) and interleukin-6 (IL-6) which stimulate the hepatic synthesis of acute phase inflammatory proteins, including CRP, which increase during malarial infection [37]. Eosinophil count was found to be elevated in both the co-infected and single infected (*Na* or *Pf*) cases compared to the negative control. Eosinophils are known to feature prominently in the leukocytic response to larval and adult stages of hookworm which is reflected by peripheral eosinophilia [38]. In general, nematode infections drive a strong Th2 response, promoting IgE synthesis [39]. Mast cell degranulation in response to IgE-allergen interaction plays a critical role in the activation and the local mobilization of eosinophils [40]. A cohort study by Kurtzhals and other [41] among children in Ghana found out that seven out of nine children with asymptomatic *P. falciparum* infection showed eosinophilia. Eosinophils have also been suggested to play a role in protection against malaria by induction of parasite killing [42]. These may account for the rise in eosinophil levels among the malaria infected subjects.

We found basophil levels to be higher in co-infected and hookworm only infected individuals compared to the malaria only or negative control. However, basophil level was higher in malaria only infected individuals than the negative control. Basophils have been poorly studied in the context of malaria with our study finding an increase in basophil count in malaria cases. It will be of paramount interest to investigate the possible mechanism accounting and the role play by basophil in malaria infection among Ghanaians.

Table 1. Characteristics of the study population recruited for the study

| Characteristic | Control (n=36) | Na (n=40) | Pf (n=59) | Na+Pf (n=63) | P-Value |
|------------------|----------------|------------------|-----------------|------------------|---------|
| Sex Male | 16 (44.4) | 17 (42.5) | 20 (33.9) | 43 (68.3) | |
| Female | 20 (55.6) | 23 (57.5) | 39 (66.1) | 20 (31.7) | |
| Mean epg (range) | 0 | 3235(144, 23328) | 0 | 2626(144, 29952) | 0.824 |
| Mean PD (range) | 0 | 0 | 1632(16, 22000) | 794(16, 12720) | 0.001 |

N. americanus (*Na*) cases, *P. falciparum* (*Pf*), *N. americanus* - *P. falciparum* co-infection (*Na+Pf*), *N. americanus* egg per gram (epg), parasite density (PD)

Table 2. Haematological parameters (mean ± SD) among study population

| Variables | Control (n=36) | Na (n=40) | Na+Pf (n=63) | Pf (n=59) | P –Value |
|------------------|---------------------------|-------------------------|-------------------------|---------------------------|----------|
| White blood cell | 5.40±1.3 | 4.85±1.7 ^a | 5.31±2.4 | 5.93± 1.5 ^a | 0.056 |
| Lymphocyte | 51.06±8.1 ^{bd} | 64.69±9.3 ^{cd} | 66.07±8.1 ^{ab} | 53.19±10.4 ^{ac} | <0.001 |
| Neutrophil | 34.29±10.1 ^{bde} | 21.48±8.9 ^{cd} | 17.50±9.8 ^{ab} | 27.09±11.3 ^{ace} | <0.001 |
| Monocyte | 9.23±2.4 ^{bd} | 4.17±2.7 ^{cd} | 5.25±3.9 ^{ab} | 10.27±3.3 ^{ac} | <0.001 |
| Eosinophil | 4.50±4.2 ^a | 6.57±6.3 | 8.22±5.8 ^a | 7.91±6.1 | 0.031 |
| Basophil | 0.93±0.3 ^{bd} | 3.09±1.8 ^{cd} | 2.96±1.6 ^{ab} | 1.55±1.9 ^{ac} | 0.01 |
| Red blood cell | 4.82±0.8 ^{ab} | 4.36±0.7 | 4.36±0.6 ^a | 4.31±0.9 ^b | 0.022 |
| Haemoglobin | 13.61±2.4 ^{ab} | 12.67±2.3 | 12.13±1.6 ^{ab} | 11.76±2.5 ^{ab} | 0.003 |
| Platelet | 220.9±92.2 | 195.3±83.9 | 184.8±65.8 | 203.9±89.7 | 0.251 |

P-values were calculated using ANOVA. Means that share a common letter are significantly different (Using the Tukey multiple comparisons test); N. americanus (Na); P. falciparum (Pf); N. americanus-P. falciparum co-infection (Na+Pf); Negative endemic control (control)

Table 3. Spearman's rank correlation coefficients for laboratory parameters and intensity of infection

| Variables | Na intensity/epg | | Pf intensity/Density | |
|-------------------|------------------|------------------|----------------------|------------------|
| | <i>r</i> | <i>P</i> | <i>r</i> | <i>P</i> |
| White blood cells | -0.235 | 0.001 | 0.080 | 0.277 |
| Lymphocytes | 0.534 | <0.001 | -0.020 | 0.785 |
| Neutrophil | -0.437 | <0.001 | -0.103 | 0.163 |
| Monocytes | -0.562 | <0.001 | 0.281 | <0.001 |
| Eosinophils | 0.103 | 0.161 | 0.154 | 0.036 |
| Basophils | 0.614 | <0.001 | -0.017 | 0.815 |
| Haemoglobin | -0.052 | 0.481 | -0.239 | 0.001 |

r, Spearman's rank coefficient; Na, N. americanus; Pf, P. falciparum

The function of eosinophilia against *N. americanus* infections in the present study population remains unclear as no significant reduction in intensity of hookworm infection was observed with an increased in eosinophil count. This is surprising, since, eosinophilia play a vital role in keeping the intensities of STH infections low by killing incoming larval stages [29, 43]. Our study finding strongly indicated that *P. falciparum* infections induce eosinophilia among Ghanaians and confirms previous study [44]. Hence, further investigation is needed to elucidate the possible protective or pathological role of eosinophil in malaria among Ghanaians.

5. CONCLUSION

The study shows that hookworm-malaria co-infection is associated with decrease in neutrophil and monocyte level but an increased in haemoglobin, eosinophil count and lymphocyte levels compared to malaria only infection. The possible mechanisms accounting for the variations in the haematological levels remain to

be elucidated and could potentially have implications on control strategies in areas where both infections are endemic.

CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

As per international standard or university standard, written approval of Ethics committee has been collected and preserved by the authors.

FUNDING

This work was supported by National Institutes of Health, grant ID #: 1R01AI099623 award MDW. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: Ascariasis, trichuriasis, and hookworm. *The Lancet*. 2006;367(9521):1521-1532.
2. Crompton DWT, Nesheim M. Nutritional impact of intestinal helminthiasis during the human life cycle. *Annu Rev Nutr*. 2002; 22(1):35-59.
3. De Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: Updating the global picture. *Trends Parasitol*. 2003; 19(12):547-551.
4. Yelifari L, Bloch P, Magnussen P, Lieshout LV, Dery G, Anemana S, et al. Distribution of human *Oesophagostomum bifurcum*, hookworm and *Strongyloides stercoralis* infections in Northern Ghana. *Trans Royal Soc Trop Med Hyg*. 2005;99(1):32-38.
5. Humphries D, Mosites E, Otchere J, Twum WA, Woo L, Jones-Sanpei H, et al. Epidemiology of hookworm infection in Kintampo North Municipality, Ghana: patterns of malaria coinfection, anemia, and albendazole treatment failure. *Am J Trop Med Hyg*. 2011;84(5):792-800.
6. Ndyomugenyi R, Kabatereine N, Olsen A, Magnussen P. Malaria and hookworm infections in relation to haemoglobin and serum ferritin levels in pregnancy in Masindi district, western Uganda. *Trans Royal Soc Trop Med Hyg*. 2008;102(2): 130-136.
7. Dickson R, Awasthi S, Williamson P, Demellweek C, Garner P. Effects of treatment for intestinal helminth infection on growth and cognitive performance in children: Systematic review of randomised trials. *Bmj*. 2000;320(7251):1697-1701.
8. Stoltzfus RJ, Kvalsvig JD, Chwaya HM, Montresor A, Albonico M, Tielsch JM, et al. Effects of iron supplementation and anthelmintic treatment on motor and language development of preschool children in Zanzibar: Double blind, placebo controlled study. *BMJ*. 2001;323(7326): 1389.
9. Dev V, Hira C, Rajkhowa M. Malaria-attributable morbidity in Assam, north-eastern India. *Annals of Tropical Medicine and Parasitology*. 2001;95(8):789-796.
10. Alonso PL, Brown G, Arevalo-Herrera M, Binka F, Chitnis C, Collins F, et al. A research agenda to underpin malaria eradication. *PLoS medicine*. 2001;8(1): e1000406.
11. Bhattarai A, Ali AS, Kachur SP, Mårtensson A, Abbas AK, Khatib R, et al. Impact of artemisinin-based combination therapy and insecticide-treated nets on malaria burden in Zanzibar. *PLoS Medicine*. 2007;4(11):e309.
12. Buck A, Anderson R, MacRae A. Epidemiology of poly-parasitism. I. Occurrence, frequency and distribution of multiple infections in rural communities in Chad, Peru, Afghanistan, and Zaire. *Tropenmedizin und Parasitologie*. 1978; 29(1):61-70.
13. World Health Organization. Soil-transmitted helminth infections. 20 February 2018. Available:<https://www.who.int/news-room/fact-sheets/detail/soil-transmitted-helminth-infections>. Assessed on 19th December, 2018
14. Henry JM, Rick MM. Helminth infections and host immune regulation. *Clin Microbiology Rev*. 2002;25(4):585-608.
15. WHO, Basic Malaria Microscopy: Part I Learner's Guide, World Health Organization, Geneva, Switzerland; 1991.
16. Monti J, Chilton N, Bao-Zhen Q, Gasser R. Specific amplification of *Necator americanus* or *Ancylostoma duodenale* DNA by PCR using markers in ITS-1 rDNA, and its implications. *Mol Cell Probe*. 1998;12(2):71-78.
17. Snounou G. Genotyping of *Plasmodium* spp. nested PCR. *Malaria Methods and Protocols: Methods and Protocols*. 2002; 103-116.
18. Padley D, Moody A, Chiodini P, Saldanha J. Use of a rapid, single-round, multiplex PCR to detect malarial parasites and identify the species present. *Ann Trop Med Parasitol*. 2003;97(2):131-137.
19. Shrivastava S, Singh N, Nigam AK, Chandel SS, Shrivastava R, Kumar S. Comparative study of hematological parameters along with effect of chemotherapy and radiotherapy in different

- stages of breast cancer. *Int J Res Med Sci.* 2017;5:311-5.
20. Brooker S, Akhwale W, Pullan R, Estambale B, Clarke SE, Snow RW, et al. Epidemiology of Plasmodium-helminth co-infection in Africa: Populations at risk, potential impact on anemia, and prospects for combining control. *Am J Trop Med Hyg.* 2007;77(6 Suppl):88-98.
 21. Brooker S, Hotez PJ, Bundy DA. Hookworm-related anaemia among pregnant women: A systematic review. *PLoS Negl Trop Dis.* 2008;2(9):e291.
 22. Chang K-H, Stevenson MM. Malarial anaemia: mechanisms and implications of insufficient erythropoiesis during blood-stage malaria. *Int J Parasitol.* 2004;34(13): 1501-1516.
 23. Price RN, Simpson JA, Nosten F, Luxemburger C, Hkirkjaroen L, ter Kuile F, et al. Factors contributing to anemia after uncomplicated falciparum malaria. *Am J Trop Med Hyg.* 2001;65(5):614-622.
 24. Ekejindu I, Okeke E, Akah B, Okpala E, Ezeagwuna D, Onwurah O. Malaria and hookworm co-infection among pregnant and non-pregnant women in a semi-urban area in Anambra State, Nigeria. *Journal of Medical Sciences.* 2001;6(1):33-35.
 25. Brooker S, Clements AC, Hotez PJ, Hay SI, Tatem AJ, Bundy DA, et al. The co-distribution of *Plasmodium falciparum* and hookworm among African schoolchildren. *Malar J.* 2006;5(1):99.
 26. Kung'u JK, Goodman D, Haji HJ, Ramsan M, Wright VJ, Bickle QD, et al. Early helminth infections are inversely related to anemia, malnutrition, and malaria and are not associated with inflammation in 6-to 23-month-old Zanzibari children. *Am J Trop Med Hyg.* 2009;81(6):1062-1070.
 27. Melo GC, Reyes-Lecca RC, Vitor-Silva S, Monteiro WM, Martins M, Benzecry SG, et al. Concurrent helminthic infection protects schoolchildren with *Plasmodium vivax* from anemia. *PLoS One.* 2010;5(6):e11206.
 28. McDevitt MA, Xie J, Gordeuk V, Bucala R. The anemia of malaria infection: Role of inflammatory cytokines. *Curr Hematol Rep.* 2004;3(2):97-106.
 29. Nacher M. Interactions between worm infections and malaria. *Clin Rev Allergy Immunol.* 2004;26(2):85-92.
 30. Nacher M, Gay F, Singhasivanon P, Krudsood S, Treeprasertsuk S, Mazier D, et al. *Ascaris lumbricoides* infection is associated with protection from cerebral malaria. *Parasite Immunol.* 2000;22(3): 107-113.
 31. Bower MA, Constant SL, Mendez S. *Necator americanus*: the Na-ASP-2 protein secreted by the infective larvae induces neutrophil recruitment in vivo and in vitro. *Exp Parasitol.* 2008;118(4):569-75.
 32. Loukas A, Prociw P. Immune responses in hookworm infections. *Clin Microbiology Rev.* 2001;14(4):689-703.
 33. Ranjit N, Jones MK, Stenzel DJ, Gasser RB, Loukas A. A survey of the intestinal transcriptomes of the hookworms, *Necator americanus* and *Ancylostoma caninum*, using tissues isolated by laser microdissection microscopy. *Int J Parasitol.* 2006;36:701-710.
 34. Eldridge L. Neutrophils function and abnormal results. Available: <https://www.verywellhealth.com/what-are-neutrophils-p2-2249134>. 2018
 35. Maina RN, Walsh D, Gaddy C, Hongo G, Waitumbi J, Otieno L, et al. Impact of *Plasmodium falciparum* infection on haematological parameters in children living in Western Kenya. *Malar J.* 2010;9(Suppl 3):S4.
 36. Abdalla SH. Peripheral blood and bone marrow leucocytes in Gambian children with malaria: numerical changes and evaluation of phagocytosis. *Ann Trop Paed.* 1988;8:250-8.
 37. Wickramasinghe SN, Abdalla SH. Blood and bone marrow changes in malaria. *Baillieres Best Pract Res Clin Haematol.* 2000;13:277-299.
 38. Allen JE, Maizels RM. Immunology of human helminth infection. *Int Arch Allergy Immunol.* 1996;109:3-10.
 39. Else KJ, Finkelman FD. Intestinal nematode parasites, cytokines and effector mechanisms. *Int J Parasitol.* 1998;28: 1145-1158.
 40. Davis TME, Ho M, Supanaranond W, Looareesuwan S, Pukrittayamee S, White NJ. Changes in the peripheral blood eosinophil count in falciparum malaria. *Acta Trop.* 1991;48:243-5.
 41. Kurtzhals JAL, Rodrigues O, Addae M, Commey JOO, Nkrumah FK, Hviid L. Reversible suppression of bone marrow response to erythropoietin in *Plasmodium*

- falciparum* malaria. Br J Haematol. 1997;97:169-74.
42. Waters LS, Taverne J, Tai P-C, Spry CJF, Targett GAT, Playfair JHL. Killing of *Plasmodium falciparum* by eosinophil secretory products. Infect Immun. 1987;55: 877-81.
43. Maizels RM, Balic A. Resistance to helminth infection: The case for interleukin-5-dependent mechanisms. Journal of Infectious Diseases 2004;190(3):427-429.
44. Kurtzhals J, Reimert C, Tette E, Dunyo S, Koram K, Akanmori B, et al. Increased eosinophil activity in acute *Plasmodium falciparum* infection-association with cerebral malaria. Clinical and Experimental Immunology. 1998;112(2):303-307.

© 2018 Amoani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47535>